

SCIENTIFIC REPORTS

OPEN

Inhibition of airway surface fluid absorption by cholinergic stimulation

Nam Soo Joo¹, Mauri E. Krouse^{1,†}, Jae Young Choi^{1,2}, Hyung-Ju Cho^{1,2} & Jeffrey J. Wine¹

Received: 15 April 2015

Accepted: 07 January 2016

Published: 05 February 2016

In upper airways airway surface liquid (ASL) depth and clearance rates are both increased by fluid secretion. Secretion is opposed by fluid absorption, mainly via the epithelial sodium channel, ENaC. In static systems, increased fluid depth activates ENaC and decreased depth inhibits it, suggesting that secretion indirectly activates ENaC to reduce ASL depth. We propose an alternate mechanism in which cholinergic input, which causes copious airway gland secretion, also inhibits ENaC-mediated absorption. The conjoint action accelerates clearance, and the increased transport of mucus out of the airways restores ASL depth while cleansing the airways. We were intrigued by early reports of cholinergic inhibition of absorption by airways in some species. To reinvestigate this phenomenon, we studied inward short-circuit currents (Isc) in tracheal mucosa from human, sheep, pig, ferret, and rabbit and in two types of cultured cells. Basal Isc was inhibited 20–70% by the ENaC inhibitor, benzamil. Long-lasting inhibition of ENaC-dependent Isc was also produced by basolateral carbachol in all preparations except rabbit and the H441 cell line. Atropine inhibition produced a slow recovery or prevented inhibition if added before carbachol. The mechanism for inhibition was not determined and is most likely multi-factorial. However, its physiological significance is expected to be increased mucus clearance rates in cholinergically stimulated airways.

In human airways, mucociliary and cough clearance are critical components of mucosal innate defense^{1,2}. Mucus clearance requires an adequate fluid layer lining the airways, and this is maintained by a balance of electrogenic, anion-mediated fluid secretion and Na⁺-mediated fluid absorption, the latter mediated mainly via the epithelial sodium channel, ENaC^{1,3}. A fundamental question in airway physiology is how airways orchestrate these opposing forces. This question has mainly been explored in cultured surface epithelia, where it has been shown that cells respond to changes in the fluid volume covering them, resulting in a homeostatically controlled depth^{4,5}. However, in upper airways, submucosal glands provide an additional and powerful component of secretion to the airways⁶, and in all airways, the rate of mucus transport provides another mechanism for adjusting the depth of airway surface liquid (ASL). In this paper, we consider what occurs when airways are stimulated cholinergically, and reconsider early experiments that showed dual actions of cholinergic stimuli on airway epithelia. We propose that in addition to promoting gland secretion, cholinergic inhibition of surface epithelial sodium absorption promotes increased surface fluid volume from the epithelia themselves. Because increased ASL depth leads to increased mucus transport rates^{7,8}, these coordinated secretory mechanisms will enhance clearance. Thus, increased mucus clearance rate is an additional homeostatic mechanism to restore ASL depth in addition concomitant with its main function of cleansing the airways.

In Ussing chamber experiments with short-circuited epithelia, the Isc across unstimulated airways of most species is dominated by sodium absorption through ENaC, so that large drops in Isc occur when ENaC is blocked. When studying the effects of agonists such as acetylcholine (ACh), most investigators first block ENaC, allowing them to focus on anion secretion. However, Na⁺ absorption by at least some airway surface epithelia is also affected by ACh. Al-Bazzaz and colleagues^{9,10}, showed a biphasic response to basolateral carbachol in sheep bronchioles. They observed an initial transient Isc increase followed by a long-lasting decrease due to inhibition of Na⁺ transport. They suggested that ACh-induced inhibition of Na⁺ absorption resulted from increased [Ca²⁺]_i, based on their prior work measuring Isc and ion fluxes across dog trachea, where it was shown that the Ca²⁺ ionophore, A23187, abolished Na⁺ transport in a Ca²⁺-dependent fashion¹¹. In excised human bronchi, 0.1 mM ACh caused

¹The Cystic Fibrosis Research Laboratory, Stanford University, Stanford, CA 94305-2130, USA. ²Department of Otorhinolaryngology, Yonsei University, Seoul, Korea. [†]Present address: Flatley Discovery lab, Charlestown, MA 02129, USA. Correspondence and requests for materials should be addressed to N.S.J. (email: nsjoo@stanford.edu)

an I_{sc} decrease, and net Na^+ absorption was abolished¹². ACh also induced an electrically silent flow of Na^+ and Cl^- , which they inferred was attributed to submucosal gland secretion. Inhibition of ENaC-mediated I_{sc} by increased $[Ca^{2+}]_i$ was subsequently shown in cultured mouse endometrial cells, where ATP was used to elevate $[Ca^{2+}]_i$ ¹³. Airway ciliated cells in humans¹⁴ and sheep contain muscarinic receptors¹⁵ (and references therein), which raise $[Ca^{2+}]_i$ and increase ciliary beat frequency when stimulated.

Airway ENaC activity is influenced in multiple ways, as expected for the key channel that gates the large electro-chemical driving force for Na^+ influx into the cell, with resulting profound effects on transmembrane ion and fluid transport. Indeed, the hyperpolarization that occurs when ENaC closes is an efficient way to induce anion secretion, because the recycling of Na^+ requires 4 times more energy than the recycling of Cl^- ¹⁶. Accordingly, factors reported to alter ENaC activity include multiple components of the cytosol^{17,18} and of the airway surface liquid (ASL), including proteases and anti-proteases^{19–21}. Importantly, many of the regulatory factors in ASL are mainly provided by the glands²².

Although inhibition of ENaC is the most obvious way to inhibit absorption, there is evidence for other mechanisms. In sheep trachea, Acevedo²³ observed biphasic I_{sc} responses to ACh with properties like those seen previously in bronchioles. Surprisingly, however, a biphasic I_{sc} response with corresponding increases and decreases in conductance was still observed after permeabilizing the apical membrane with amphotericin and imposing a K^+ gradient, suggesting that ACh was transiently activating and then inhibiting basolateral K^+ channels under these conditions.

With this background, we re-examined cholinergic effects on airway mucosal I_{sc} with a view to understanding its generality and physiological significance.

Results

Carbachol inhibited ENaC-dependent I_{sc} stimulation in four species. I_{sc} was measured across excised tracheal epithelia of 5 species: humans, sheep, pigs, ferrets, and rabbits. I_{sc} was also measured across planar sheets of cultured primary human bronchial epithelia (HBE) and H441 human small airway cells. The basic procedure was to measure resting I_{sc} and then the I_{sc} response to carbachol added either before or after benzamil. In all tissues except ferret tracheas the resting I_{sc} was inhibited 50–90% by benzamil; in ferrets benzamil inhibited <20% of the resting I_{sc} .

In sheep, 100 μ M basolateral carbachol stimulated a transient increase ($t_{1/2} = 88.9 \pm 10.6$ sec, $n = 15$) in I_{sc} followed by a sustained decrease ($t_{1/2} > 30$ min, Fig. 1A, *bottom I_{sc} trace*). When conditions were reversed and benzamil was added first, it produced a sustained decrease in I_{sc} as expected, and addition of carbachol now stimulated only the transient increase in I_{sc} ; additional benzamil was without effect (Fig. 1A, *top I_{sc} trace*). Results for sheep are summarized in Fig. 1B and Supplementary Table S.1. Carbachol inhibited the benzamil-sensitive (ENaC-dependent) I_{sc} by $80 \pm 3\%$ ($n = 15$, 13 sheep, $P < 0.0001$, Fig. 1B), whereas carbachol given after benzamil produced an initial transient increase ($167 \pm 52\%$, $n = 3$ sheep, $P = 0.015$) followed by a sustained increase in I_{sc} ($38 \pm 29\%$, $n = 3$ sheep, n.s. $P = 0.14$) (Fig. 1B). Neither of these increases were sufficient to restore the original I_{sc} .

Basolateral carbachol also inhibited ENaC-dependent (benzamil-sensitive) I_{sc} in pig, human (Fig. 1B), some ferret tracheas (Supplementary Figure 1C) and in primary HBE cultures (see below). Apical carbachol had no consistent effect in any of these preparations (Supplementary Figure 1B). Carbachol inhibition of sheep and pig airways occurred with concentrations as low as 1 μ M, but inhibition at that concentration developed slowly, making it difficult to study. In sheep, 100 μ M carbachol rapidly and reliably induced inhibition; in the other species 1 mM carbachol was required to consistently inhibit ENaC-dependent I_{sc} , and in ferrets even that concentration gave inconsistent results. (Note that ferret experiments were done in the presence of cartilage, and ferrets also had the smallest response to benzamil, see below).

In human large airways $76.7 \pm 11.7\%$ of the I_{sc} was benzamil-sensitive, and carbachol inhibited $39 \pm 3.3\%$ of that I_{sc} ($n = 4$ human subjects). In pig tracheas $58.8 \pm 2.6\%$ of the I_{sc} was ENaC-dependent and carbachol inhibited $44.2 \pm 4.7\%$ of that I_{sc} ($n = 10$ pigs). In ferret tracheas only $17.3 \pm 4.2\%$ of I_{sc} was inhibited by benzamil ($n = 32$, 16 ferrets). When carbachol was added first, it inhibited the I_{sc} in only 3 of 8 ferrets by $36 \pm 20.5\%$ (Supplementary Figure 1C).

Carbachol inhibited ENaC-dependent I_{sc} in primary cultures of HBE cells. Stimulation of HBE cells with 1 mM carbachol produced an I_{sc} drop in all monolayer HBE cell inserts ($n = 6$, Figs 1B and 2 and Supplementary Table S.1). The initial transient increase in I_{sc} was lacking or small. (The transient increase in human airways was also small, see below). In contrast with explanted airways, benzamil subsequently caused a second large drop in I_{sc} (*top I_{sc} trace*) to $\sim$ zero I_{sc} , and when benzamil was added first it almost abolished the I_{sc} ($87.4 \pm 6.9\%$, $n = 3$), again in contrast with explanted airway tissues, where $\sim 30\%$ of the original I_{sc} remained after benzamil. I_{sc} recovered slowly after wash out to 60–70% of the I_{sc} which could be inhibited with repeated benzamil (Fig. 2, *top I_{sc} trace*). If benzamil was added first (Fig. 2, *bottom I_{sc} trace*), the subsequent washout increased I_{sc} and part of the recovered I_{sc} was sensitive to 1 mM carbachol. To our knowledge this is the first demonstration of carbachol-induced inhibition of primary HBE cells.

Inhibition was unchanged after reducing anion transport. The decline in I_{sc} following carbachol might partially result from inhibition of anion secretion, although that interpretation is not consistent with the small increases in I_{sc} seen when carbachol was added after benzamil. To further explore a role for anion secretion, we stimulated sheep trachea with carbachol in normal Krebs-Ringer bicarbonate (KRB) buffer and in a buffer designed to block anion transport (HEPES for replacing HCO_3^- and 100 μ M bumetanide to block, $Na^+/K^+/2Cl^-$ cotransporter, NKCC). These conditions reduce carbachol-stimulated gland secretion by $\sim 90\%$ ²⁴. No significant difference in carbachol-induced inhibition of the ENaC-dependent I_{sc} was observed when anion




Figure 1. Carbachol inhibits Na^+ transport across airway surface epithelia. (A,B) Ussing chamber I_{sc} responses (1 mV pulse every 20 seconds) from a matched pair of sheep tracheal mucosa to drug additions indicated by arrows. (A) Carbachol (100 μM , basolateral, 'Carb') followed by 10 μM apical benzamil (Bz) (*top I_{sc} trace*). Drugs given in reverse order: carbachol following benzamil (*bottom I_{sc} trace*). (B) Summary of carbachol-induced ENaC inhibition in airways from three different species ($n = 4, 13$, and 10 human, sheep, and pig, respectively) and primary human bronchial epithelial cells (HBE, $n = 9$). Carbachol concentration was 1 mM except for sheep (100 μM). Bz-first represents the benzamil treatment preceded carbachol treatment and the bars are showing peak of the transient increase (open bar), sustained (black bar) I_{sc} responses-induced by carbachol, and I_{sc} response-inhibited by benzamil (grey bar).




Figure 2. Carbachol inhibition of I_{sc} in primary cultures of human bronchial epithelial cells. Top trace: ENaC-dependent I_{sc} is partially inhibited by 1 mM carbachol and reduced to zero with 10 μM benzamil. Both kinds of inhibition were partially reversed by washing. Bottom trace: all I_{sc} was inhibited by benzamil but not by carbachol. Unlike animal tracheas, an initial transient I_{sc} increase to 1 mM carbachol was either small or absent in HBE cells ($n = 9$ monolayer inserts).

transport was reduced by bumetanide + HEPES (the residual ENaC-dependent ENaC I_{sc} : 11.6 ± 2.7 in KRB and $11.4 \pm 2.6 \mu A/cm^2$, $P = 0.8$, $n = 6$ from 4 sheep), while the transient I_{sc} increase was greatly reduced in HEPES + bumetanide buffer (14.7 ± 2.4 in KRB and $7.8 \pm 2.8 \mu A/cm^2$ in HEPES + Bm) (Figure 3).

Carbachol inhibition of I_{sc} is mediated by muscarinic receptors. Carbachol, like acetylcholine, stimulates both muscarinic and nicotinic receptors. Furthermore, the high concentrations needed to see inhibition




Figure 3. Na^+ absorption but not anion secretion was inhibited by cholinergic stimulation. (A) (top trace, anion transport inhibitors present, bottom trace, control buffer) In sheep trachea, ENaC-dependent I_{sc} inhibition by carbachol persisted after anion secretion was blocked by 100 μM basolateral bumetanide (Bm) plus HCO_3^- replacement by HEPES with pure O_2 supply. The transient I_{sc} increase by carbachol was greatly reduced in the bumetanide and HEPES condition. (B) Average magnitude of residual benzamil-sensitive I_{sc} in sheep airways after carbachol; control (KRB, Krebs-bicarbonate) vs. HEPES/Bm buffers ($n = 6$ each, from 4 sheep).

of I_{sc} in some preparations raised a concern about off-target effects. To determine if I_{sc} inhibition was mediated by muscarinic receptors, we compared effects of the muscarinic antagonist atropine and the nicotinic antagonist hexamethonium bromide on carbachol inhibition of ENaC-dependent I_{sc} . When atropine was added near the maximum of carbachol-induced inhibition it led to recovery with a time course similar to that seen after wash-out of carbachol (Fig. 4A). When atropine (1 or 10 μM) was added basolaterally before carbachol, it abolished the I_{sc} decreases to carbachol (Fig. 4B) ($n = 4$ sheep and 2 pigs). In contrast, hexamethonium bromide up to 100 μM had no effect on carbachol-induced inhibition (Supplementary Figure 2A), and nicotine (1 mM, basolateral) failed to alter I_{sc} in pig or sheep tracheal mucosa (3 pigs and 2 sheep, Supplementary Figure 2B).

Apical ATP inhibited ENaC-dependent I_{sc} , but was not required for carbachol-induced inhibition. In several systems^{13,18,25,26}, apical ATP inhibits amiloride-sensitive I_{sc} . To investigate the possibility that carbachol-induced inhibition might be mediated by release of ATP, sheep tracheas were pretreated with 100 μM ATP before carbachol addition. As previously described in other systems, ATP decreased the ENaC-dependent I_{sc} . However, subsequent addition of 100 μM carbachol produced an additional decrease in I_{sc} (Fig. 5A, top I_{sc} trace). When ATP was added after carbachol-induced inhibition had plateaued, it produced no further inhibition (Fig. 5A, bottom I_{sc} trace). Summary data (Fig. 5B) show that ATP alone reduced ENaC-dependent I_{sc} by $48.4 \pm 8.1\%$ ($n = 3$ sheep, $P = 0.016$) and subsequent carbachol addition almost abolished the remaining ENaC-dependent I_{sc} ($P = 0.001$). These results show that carbachol inhibits ENaC-dependent I_{sc} more effectively than does ATP, but do not eliminate the possibility that ATP release is a component of carbachol-induced inhibition. To evaluate that possibility we added 10 U/ml apical apyrase (which catalyzes the hydrolysis of ATP), to sheep and pig airway mucosa ($n = 7$ from 4 sheep and one pig) prior to adding carbachol or ATP. Apyrase pretreatment abolished ATP-induced inhibition but had no effect on the carbachol-induced inhibition (Figure 5C and D). Thus, if ATP is released into the ASL by carbachol its inhibitory effects are occlusive with other inhibitory mechanisms.

Carbachol failed to inhibit I_{sc} in rabbit trachea and in the human H441 cell line. In rabbit tracheas ~50% of total I_{sc} was benzamil-sensitive (ENaC-dependent) prior to stimulation ($n = 3$ rabbits). When rabbit tracheal mucosa was treated with 100 μM carbachol either to basolateral or apical side before or after benzamil, it produced no change in the I_{sc} (Fig. 6A, $n = 4$ from 3 rabbits). This lack of responsiveness by rabbit tracheas is consistent with an earlier report that studied excised tracheas²⁷. However, cultured rabbit tracheal cells responded to carbachol with increased I_{sc} and $[Ca^{2+}]_i$ ²⁸. We confirmed that rabbit tracheal I_{sc} was inhibited by apical ATP (Supplementary Figure 3A), as first shown by Iwase and colleagues²⁵.

H441 cells are human small airway cells that express functional ENaC. H441 cells developed robust ENaC-dependent I_{sc} (~93% of their starting I_{sc} : 38.5 ± 4.5 to $2.8 \pm 0.8 \mu A/cm^2$, $n = 7$ inserts) but they did not respond to basolateral or apical application of 1 mM carbachol (Fig. 6B, $n = 6$ inserts). Apical ATP produced no inhibition, instead causing a sustained increase in I_{sc} of 6 or 12 $\mu A/cm^2$ ($n = 2$ inserts, Supplementary




Figure 4. Carbachol inhibition of I_{sc} is mediated by muscarinic receptors. (A) Carbachol ($100\mu\text{M}$) inhibition of I_{sc} in sheep trachea recovered slowly after washout (*top I_{sc} trace*) and after application of atropine (Atrp, $10\mu\text{M}$, basolateral, *bottom I_{sc} trace*). Inhibition could be repeated (*top I_{sc} trace*). (B) Dose-dependent block of cholinergic effects on I_{sc} by atropine. Sheep tracheas treated with 1 or $10\mu\text{M}$ atropine prior to 1 to $100\mu\text{M}$ carbachol. Both the initial increase and the later decrease in I_{sc} are blocked. Atropine block was observed in 6 experiments with sheep ($n = 3$) and pigs ($n = 2$, Supplementary Figure 2A).

Figure 3B, *top I_{sc} trace*). However, ENaC-dependent I_{sc} was inhibited by $4\text{--}16\mu\text{A}/\text{cm}^2$ by thapsigargin, which elevates $[\text{Ca}^{2+}]_i$. (Supplementary Figure 3B, *bottom I_{sc} trace*, $n = 2$ inserts).

These observations indicate that rabbit tracheal and H441 cells also have mechanisms for inhibiting ENaC-dependent I_{sc} , but rabbit tracheal cells either lack muscarinic receptors or fail to couple them to pathways that inhibit the ENaC-dependent I_{sc} , while H441 cells lack mechanisms to couple either muscarinic or purinergic receptors to inhibition of ENaC-dependent I_{sc} .

Proteases and anti-proteases influenced I_{sc} in H441 cells and airway mucosa. We previously proposed that airway submucosal glands could participate in the regulation of ENaC activity by secreting ENaC regulating factors, such as serine protease inhibitors, onto the airway surface when glands are stimulated^{21,29}. In our present study we found that the serine protease inhibitor aprotinin significantly reduced ENaC-dependent I_{sc} in H441 cell monolayers when $30\mu\text{g}/\text{ml}$ was applied apically (control: from 48.8 ± 3.6 to 41.2 ± 2.4 and aprotinin treated: from 47.2 ± 3.8 to $21.2 \pm 2.3\mu\text{A}/\text{cm}^2$, $P = 0.001$, $n = 4$ inserts). Trypsin partially counteracted carbachol-induced inhibition in pig and sheep airways (Supplementary Figure 3C and D). These results are consistent with our earlier interpretations^{21,29} and other reports^{19,30}.

Phospholipase-C activation was not required for carbachol-induced inhibition. A phospholipase-C pathway has been implicated in the inhibition of ENaC by flagellin³¹. To determine if cholinergic inhibition of ENaC requires PLC, we tested carbachol-induced inhibition $\pm$ the PLC inhibitor, U73122, in sheep and pig tracheas. Apical + basolateral U73122 ($10\mu\text{M}$) had no effect on carbachol inhibition of ENaC-dependent I_{sc} (Supplementary Figure 4A).

CFTR activation with forskolin did not affect carbachol inhibition of ENaC-dependent I_{sc} . CFTR interactions with ENaC have been controversial^{3,32}. To look for possible interactions, we activated or inhibited CFTR prior to carbachol stimulation and noted effects on ENaC-dependent I_{sc} . The residual ENaC-dependent I_{sc} after carbachol was not significantly different in the presence ($2.7 \pm 0.6\mu\text{A}/\text{cm}^2$) or absence ($2.8 \pm 0.5\mu\text{A}/\text{cm}^2$) of $10\mu\text{M}$ forskolin pre-treatment ($n = 5$, 4 sheep, Supplementary Figure 4B).

Discussion

Main findings. Cholinergic stimulation causes both gland secretion and increased rates of mucociliary clearance (MCC)^{7,8}. Because mucus clearance rates increase as the depth of ASL increases^{1,2}, increased MCC after cholinergic stimulation could result simply from increased ASL depth following gland secretion. This effect will




Figure 5. Apical ATP is not required for carbachol-induced inhibition of I_{sc} . (A) Inhibition of I_{sc} across sheep trachea by 100 μ M apical ATP before (top trace) and after (bottom trace) 100 μ M basolateral carbachol. Inhibition of I_{sc} by ATP is much smaller after carbachol treatment. Benzamil (10 μ M) was applied apically. (B) Summary of ATP results in sheep airways ($n = 3$); ***significantly different from control, “after,” $P < 0.05$ and $P < 0.005$, respectively. (C) In sheep trachea, apyrase pretreatment (10 U/ml, apical) had no effect on carbachol induced inhibition of ENaC-dependent I_{sc} . (D) By contrast, in sheep trachea apyrase (top trace) abolished ATP induced inhibition of ENaC-dependent I_{sc} (shown without apyrase in bottom trace). Representative traces, $n = 6$ experiments from 4 sheep.

be counterbalanced by absorption, which in cultured epithelial cells increases as fluid depth increases^{1,2}. This is a means of eventually restoring conditions to the basal state, but during the need for rapid clearance we hypothesized that the effects of glandular secretion on MCC could be amplified by switching the surface epithelia from fluid absorption to fluid secretion—indeed in one system this has been shown directly by inhibiting ENaC with benzamil⁷. How is this normally accomplished *in vivo*? The main point of this paper is that long lasting inhibition of ENaC-dependent fluid absorption is the most important factor in the switch of surface epithelia from absorption to secretion.

In the present experiments we explored the generality and physiological significance of cholinergic inhibition of ENaC-dependent I_{sc} by testing the effects of cholinergic stimulation on I_{sc} in seven types of airway preparations that display ENaC-dependent I_{sc} . Of these, five also displayed cholinergic inhibition of ENaC-dependent I_{sc} . The exceptions were rabbit trachea and the H441 human small airway cell line; we suspect but did not establish that these cells lack muscarinic receptors.

Comparison with apparently contradictory results. Some prior studies do not report cholinergic inhibition of ENaC-dependent I_{sc} . A previous study of dog tracheas found no effect of acetylcholine on inhibition of inward Na^+ current, and only a slight inhibition of Na^+ flux³³. Dogs, like rabbits, may lack this response, but the concentration of ACh was less and the measurement duration shorter than we used, so dog tracheas will need to be re-examined using comparable conditions to establish this point.

In an elegant study, results opposite to those reported here were found in cultured rat alveolar type II cells, where carbachol activated ENaC activity³⁴. Different actions of muscarinic agents in different cell types are not contradictory. Indeed, if viewed functionally, they are complementary actions to remove fluid from the airways. Fluid removal from the lung alveoli by A2 cells occurs by direct fluid absorption, which was increased by systemic carbachol³⁴. However, for the airways, with their vastly reduced surface area compared to alveoli, the mechanism for fluid removal is primarily via mucociliary clearance, which is accelerated when ENaC-dependent absorption is inhibited⁷ (Joo, unpublished observation with *ex vivo* ferret tracheas). Thus, paradoxically, inhibition of airway epithelial absorption will increase the rate of fluid (mucus) clearance from the airways to restore ASL depth. As




Figure 6. Carbachol did not inhibit ENaC-dependent I_{sc} in rabbit airways or H441 cells. (A) Rabbit tracheal preparations were treated with 100 μ M carbachol basolaterally (C-bl) and apically (C-ap) after (top trace) and before (bottom trace) 10 μ M apical benzamil (Bz) as indicated by arrows. Representative Ussing I_{sc} records from $n = 3$ rabbits. (Apical ATP (100 μ M) did inhibit ENaC-dependent I_{sc} , see Supplementary Figure 3A). (B) H441 human small airway Clara epithelial cells failed to respond to 1 mM carbachol applied to both sides of the filter (top trace), in spite of having a large ENaC-dependent I_{sc} (bottom trace). Representative traces from 6 H441 cell inserts.

mentioned above, this mechanism can't operate in closed, cell culture systems, where fluid absorption is the only means for decreasing depth.

Mechanisms of carbachol inhibition of ENaC-dependent I_{sc} : apical mediators. An efficient way to couple gland secretion to inhibition of epithelial surface absorption would be to have the glands themselves secrete factors to inhibit ENaC²² (and references therein). This hypothesis is supported by evidence that ENaC is inhibited by multiple factors in the ASL^{17–20}, many of which are abundant in glands^{6,21,22,29}. For example, the BPI fold protein SPLUNC1 inhibits ENaC activity and fluid absorption of airway surface epithelia and has been proposed as the major endogenous ENaC regulator of upper airways surface epithelia²⁰. BPI fold proteins are most highly expressed in submucosal glands in the upper airways^{35,36} and are the most abundantly detected innate defense proteins in pure mucus secreted from human airway submucosal glands²². In Ussing chambers the ASL is diluted extensively so that fluctuations in ASL soluble factors are unlikely to play a significant role in the responses we observed. However, to show that such factors can in principle participate in the cholinergic response we showed that the antiproteases, aprotinin decreased I_{sc} in H441 cells and the protease trypsin increased it in sheep and pig airway. It remains necessary to demonstrate directly that stimulating glands inhibits benzamil-sensitive I_{sc} and if so, to assess the importance of their role.

The logic is compelling that glandular secretions increase mucus clearance rates both by direct secretion of mucus and by switching the surface epithelia to a secretory mode by inhibiting ENaC-dependent fluid absorption.

Mechanisms of carbachol inhibition of ENaC-dependent I_{sc} : basolateral mediators. In spite of controversy about whether epithelia are innervated^{37–39}, human¹⁴ and sheep airway ciliated cells¹⁵ express muscarinic receptors, and 4 of the 5 species tested plus cultured HBE cells responded to the muscarinic agonist carbachol. We do not know the cellular mechanisms that intervene between activation of muscarinic receptors and inhibition of ENaC-dependent I_{sc} , but the combined results of many investigators added to our own make it clear that multiple mechanisms must be involved. It needs to be emphasized that there is no direct evidence from any study that ENaC activity of tracheal epithelial cells is decreased by cholinergic stimulation, although that is the assumption. ENaC P_o can be decreased in tracheal cells by ATP¹⁷ and increased in cultured A2 cells by muscarinic stimulation³⁴. But even in the careful analysis of A2 cells only the early links in the pathway from muscarinic receptors were described³⁴.

An alternative or additional mechanism is the inhibition of basolateral K^+ channels²³. Because the anion secretion that continues in the presence of carbachol (Fig. 1, Supplementary Figures 5A and B) is sensitive to the anion channel inhibitor, niflumic acid, this inhibition can only be partial, since some residual K^+ conductances are necessary to sustain the anion secretion. In addition to these potential mechanisms, ENaC-dependent *I*_{sc} is modulated by the membrane potential, so that cholinergic activation of anion conductances will reduce the driving force for Na^+ entry independent of any effect on ENaC number or P_O . In sum, detailed patch clamp and cell biological analyses of carbachol-induced inhibition of ENaC-dependent *I*_{sc} are warranted and feasible now that we have demonstrated the phenomenon in cultured cells HBE cells, but our focus has been on the physiological relevance, not the mechanism.

Limitations of this study. In addition to the limitations discussed above, methods beyond *I*_{sc} studies will be required to establish mechanisms. The high concentrations of carbachol that were required to see ENaC-dependent *I*_{sc} are puzzling, because it is unlikely that these occur physiologically, even as part of strong parasympathetic activation during the lung defense reflex⁴⁰. This is especially so since direct cholinergic synapses on airway epithelia have not been described and the source of ACh is likely to be more diffuse. We are confident that these are not off-target effects, because atropine eliminated the high dose carbachol effects (Supplementary Figure 2A-bottom *I*_{sc} trace). For cholinergic mediation of gland secretion, co-transmitters such as VIP produce markedly synergistic effects⁶, so it is possible that our experiments lack co-factors needed to amplify inhibition by carbachol. Also, at concentrations of carbachol below threshold for most preparations, those that responded showed a very slowly developing inhibition that eventually reached a level similar to that produced rapidly by much higher concentrations (Supplementary Figure 6). The variability and very slow development of responses to lower concentrations precluded its further study, but it remains possible that intact airways might exert tonic control of fluid absorption with low levels of acetylcholine (and perhaps with other co-transmitters).

Physiological relevance: parasympathetic coordination of optimal mucus clearance. This work was influenced by earlier work showing that ACh increased ciliary beat frequency via an M3 receptor present in airway epithelial cells¹⁵, by evidence that carbachol stimulated gland secretion and mucociliary clearance rates in pig⁷ and ferret⁸ tracheas, and that benzamil speeded MCC rates after anion transport had been diminished⁷. We hypothesize that parasympathetic stimulation *in vivo* will also inhibit ENaC-dependent, Na^+ -mediated fluid absorption, and help switch the balance of epithelia transport to net secretion, thus augmenting gland secretion and increased ciliary beat frequency to speed mucus clearance. If this interpretation is correct, it suggests caution in the use of inhaled anti-cholinergic agents to treat patients with obstructive pulmonary diseases, since these might actually slow MCC⁴¹.

Materials & Methods

Airway tissue procurement. Human tracheal scraps from donors were obtained from lung transplants harvested less than 2 h post-mortem at Stanford Hospital. The protocol for handling human airway tissues was approved and informed consent was obtained from all participants prior to the study (Stanford University-IRB protocol#: 11638). Postmortem (<1 hr) tracheas from young male and female *Yorkshire* pigs (30–50 kg), young male *Suffolk-Rambouillet* sheep (~70 kg), *New Zealand White* rabbits at the age of 5–6 months, and 3–10 months old *M. Putorius* ferrets were obtained from the animal facility at Stanford University after acute experiments unrelated to our studies. All protocols for handling animal tissues at Stanford were approved by Administrative Panel on Laboratory Animal Care (Stanford's Institutional Animal Care and Use Committee: IACUC protocol#: 10048). Methods for both human and animal tissues were carried out in accordance with approved guidelines. Harvested tissues were placed in cold PhysioSolTM (Hospira, IL) after removal for transport to the laboratory and then transferred to ice-cold Krebs Ringer bicarbonate (KRB) buffer gassed with 95% O_2 and 5% CO_2 and kept until use. The KRB buffer composition was 115 mM NaCl, 2.4 mM K_2HPO_4 , 0.4 mM KH_2PO_4 , 25 mM $NaHCO_3$, 1.2 mM $MgCl_2$, 1.2 mM $CaCl_2$, and 10 mM Glucose (pH 7.2 at room temperature) adjusted to ~290 mosM with a Wescor vapor pressure osmometer. The Krebs buffer contained 1.0 μ M indomethacin to minimize endogenously generated prostaglandins during tissue preparation. Where HCO_3^- -free bathing buffer solution was used, HCO_3^- was replaced by HEPES, 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid, and gassed with 100% O_2 . Tracheal mucosa for Ussing chamber experiments were carefully dissected from underlying cartilage while those from rabbits and ferrets were used intact without dissecting cartilage and connective tissues and only cartilaginous portions of trachea were used.

Cell culture. The human airway surface epithelial cell line²¹, H441, was maintained in a T₂₅ tissue culture flask containing a 1:1 mixture of Dulbecco's modified Eagle's medium and Ham's F-12 nutrient mixture supplemented with 10% fetal bovine serum (Sigma, MO), 100 μ g/ml streptomycin, 100 units/ml penicillin, and 2 mM glutamine at 37 °C in a humidified atmosphere containing 5% CO_2 . Cells were passaged once at 7–10 days and seeded on HPC (human placental collagen, Sigma)-coated filters (Snapwells, Corning-Costar). To provide an optimal air-interface cultures, the serum medium was switched with 2% Ultrosor-G (BioSeptra, Cergy, France), a serum substitute, containing medium in cells grown on filters 3–4 days after seeding. Human airway primary cell cultures used in this study were generously provided by Philip Karp, Jonathan Widdicombe and Walter Finkbeiner.

Electrophysiology. Human and animal airway tissue preparations and cultured cells on filters were mounted in EasyMount Ussing chambers (Physiologic Instruments, CA) with exposed surface areas of 0.07–1.13 cm² depending on the preparation, and were bathed in KRB buffer at 37 °C with a continuous 95% O_2 and 5% CO_2 gas supply, except for HCO_3^- -free buffer solutions which were gassed with pure O_2 . Transepithelial short-circuit current (*I*_{sc}) was measured using a VCC-600 voltage clamp (Physiologic Instruments, CA) and *I*_{sc} data were obtained and displayed with PowerLab Chart4 software (ADInstruments, CA). Total tissue conductance was

calculated by applying Ohm's law to the Isc deflection resulting from 1 mV pulse across the tissues/cell cultures every 20 sec during the experiment (Supplementary Table S.1).

Reagents. Chemicals were obtained from Sigma/Aldrich (St. Louise, MO) and were made fresh or maintained at -20°C as aliquots of stock concentrations. Stock solutions of carbachol, ATP, apyrase (ATP diphosphohydrolase), aprotinin, trypsin, and hexamethonium bromide were made in sterilized deionized water and bumetanide, an inhibitor of $\text{Na}^{+}/\text{K}^{+}/2\text{Cl}^{-}$ transporter in alkaline solution while benzamil, CFTRinh172, GlyH101, thapsigargin, U73122 (a phospholipase C, PLC, inhibitor), and atropine were dissolved in dimethyl sulfoxide (DMSO, Me_2SO). Carbachol-induced Isc responses were indistinguishable from those produced by acetylcholine (Supplementary Figure 1A). Therefore, we used carbachol (a metabolically non-hydrolyzable analog) instead of acetylcholine to avoid potential variability that could be caused by differing levels of acetylcholinesterase across species. Benzamil was chosen because it is a more potent inhibitor of ENaC than amiloride, giving maximal inhibition at $10\mu\text{M}$ ⁴².

Statistics. Data are presented as mean $\pm$ S.E.M. unless indicated otherwise. Student's *t*-test was used to compare means of different treatment groups. ENaC-dependent Isc was calculated either by subtracting the starting Isc from the Isc after benzamil (benzamil-first, no carbachol pretreatment) or by subtracting starting Isc from Isc after benzamil (but carbachol treatment- preceded) (Fig. 1B and Supplementary Table 1). Benzamil-sensitive Isc is alternately referred to as ENaC-dependent Isc.

References

1. Donaldson, S. H., Corcoran, T. E., Laube, B. L. & Bennett, W. D. Mucociliary clearance as an outcome measure for cystic fibrosis clinical research. *Proceedings of the American Thoracic Society* **4**, 399–405 (2007).
2. Knowles, M. R. & Boucher, R. C. Mucus clearance as a primary innate defense mechanism for mammalian airways. *Journal of Clinical Investigation* **109**, 571–577 (2002).
3. Itani, O. A. *et al.* Human cystic fibrosis airway epithelia have reduced Cl^{-} conductance but not increased Na^{+} conductance. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 10260–10265 (2011).
4. Matsui, H., Davis, C. W., Tarran, R. & Boucher, R. C. Osmotic water permeabilities of cultured, well-differentiated normal and cystic fibrosis airway epithelia. *J Clin Invest* **105**, 1419–1427 (2000).
5. Tarran, R., Grubb, B. R., Gatzky, J. T., Davis, C. W. & Boucher, R. C. The relative roles of passive surface forces and active ion transport in the modulation of airway surface liquid volume and composition. *The Journal of general physiology* **118**, 223–236 (2001).
6. Widdicombe, J. H. & Wine, J. J. Airway Gland Structure and Function. *Physiol Rev* **95**, 1241–1319, 2014 (2015).
7. Ballard, S. T., Trout, L., Mehta, A. & Inglis, S. K. Liquid secretion inhibitors reduce mucociliary transport in glandular airways. *Am J Physiol Lung Cell Mol Physiol* **283**, L329–335, 2001 (2002).
8. Jeong, J. H., Joo, N. S., Hwang, P. H. & Wine, J. J. Mucociliary clearance and submucosal gland secretion in the *ex vivo* ferret trachea. *Am J Physiol Lung Cell Mol Physiol* **307**, L83–93 (2014).
9. Al-Bazzaz, F. J. Regulation of Na and Cl transport in sheep distal airways. *Am J Physiol* **267**, L193–198 (1994).
10. Al-Bazzaz, F. J. & Gailey, C. Ion transport by sheep distal airways in a miniature chamber. *Am J Physiol Lung Cell Mol Physiol* **281**, L1028–1034 (2001).
11. Al-Bazzaz, F. & Jayaram, T. Ion transport by canine tracheal mucosa: effect of elevation of cellular calcium. *Experimental lung research* **2**, 121–130 (1981).
12. Knowles, M. *et al.* Bioelectric properties and ion flow across excised human bronchi. *J Appl Physiol Respir Environ Exerc Physiol* **56**, 868–877 (1984).
13. Wang, X. F. & Chan, H. C. Adenosine triphosphate induces inhibition of Na^{+} absorption in mouse endometrial epithelium: a Ca^{2+} -dependent mechanism. *Biology of reproduction* **63**, 1918–1924 (2000).
14. Corssen, G. & Allen, C. R. Acetylcholine: its significance in controlling ciliary activity of human respiratory epithelium *in vitro*. *J Appl Physiol* **14**, 901–904 (1959).
15. Salathe, M., Lipson, E. J. & Ivonnet, P. I. & Bookman, R. J. Muscarinic signaling in ciliated tracheal epithelial cells: dual effects on Ca^{2+} and ciliary beating. *The American journal of physiology* **272**, L301–310 (1997).
16. Welsh, M. J. Energetics of chloride secretion in canine tracheal epithelium. Comparison of the metabolic cost of chloride transport with the metabolic cost of sodium transport. *Journal of Clinical Investigation* **74**, 262–268 (1984).
17. Kunzelmann, K. *et al.* Purinergic inhibition of the epithelial Na^{+} transport via hydrolysis of PIP₂. *Faseb J* **19**, 142–143 (2005).
18. Tarran, R., Trout, L., Donaldson, S. H. & Boucher, R. C. Soluble mediators, not cilia, determine airway surface liquid volume in normal and cystic fibrosis superficial airway epithelia. *J Gen Physiol* **127**, 591–604 (2006).
19. Caldwell, R. A., Boucher, R. C. & Stutts, M. J. Neutrophil elastase activates near-silent epithelial Na^{+} channels and increases airway epithelial Na^{+} transport. *Am J Physiol Lung Cell Mol Physiol* **288**, L813–819 (2005).
20. Garcia-Caballero, A. *et al.* SPLUNC1 regulates airway surface liquid volume by protecting ENaC from proteolytic cleavage. *Proc Natl Acad Sci USA* **106**, 11412–11417 (2009).
21. Joo, N. S., Irokawa, T., Robbins, R. C. & Wine, J. J. Hyposecretion, not hyperabsorption, is the basic defect of cystic fibrosis airway glands. *J Biol Chem* **281**, 7392–7398 (2006).
22. Joo, N. S. *et al.* Proteomic analysis of pure human airway gland mucus reveals a large component of protective proteins. *PLoS ONE* **10**, e0116756 (2015).
23. Acevedo, M. Effect of acetyl choline on ion transport in sheep tracheal epithelium. *Pflugers Arch* **427**, 543–546 (1994).
24. Joo, N. S., Wu, J. V., Krouse, M. E., Saenz, Y. & Wine, J. J. Optical method for quantifying rates of mucus secretion from single submucosal glands. *Am J Physiol Lung Cell Mol Physiol* **281**, L458–468 (2001).
25. Iwase, N. *et al.* ATP-induced Cl^{-} secretion with suppressed Na^{+} absorption in rabbit tracheal epithelium. *Respiration physiology* **107**, 173–180 (1997).
26. Mason, S. J., Paradiso, A. M. & Boucher, R. C. Regulation of transepithelial ion transport and intracellular calcium by extracellular ATP in human normal and cystic fibrosis airway epithelium. *Br J Pharmacol* **103**, 1649–1656 (1991).
27. Jarnigan, F., Davis, J. D., Bromberg, P. A., Gatzky, J. T. & Boucher, R. C. Bioelectric properties and ion transport of excised rabbit trachea. *J Appl Physiol* **55**, 1884–1892 (1983).
28. Van Scott, M. R. & Paradiso, A. M. Intracellular Ca^{2+} and regulation of ion transport across rabbit Clara cells. *Am J Physiol* **263**, L122–127 (1992).
29. Joo, N. S., Lee, D. J., Wings, K. M., Rustagi, A. & Wine, J. J. Regulation of antiprotease and antimicrobial protein secretion by airway submucosal gland serous cells. *J Biol Chem* **279**, 38854–38860 (2004).
30. Myerburg, M. M. *et al.* Airway surface liquid volume regulates ENaC by altering the serine protease-protease inhibitor balance: a mechanism for sodium hyperabsorption in cystic fibrosis. *J Biol Chem* **281**, 27942–27949 (2006).
31. Kunzelmann, K. *et al.* Flagellin of *Pseudomonas aeruginosa* inhibits Na^{+} transport in airway epithelia. *FASEB J* **20**, 545–546 (2006).

32. Stutts, M. J. *et al.* CFTR as a cAMP-dependent regulator of sodium channels. *Science* **269**, 847–850 (1995).
33. Boucher, R. C. & Gatz, J. T. Regional effects of autonomic agents on ion transport across excised canine airways. *J Appl Physiol* **52**, 893–901 (1982).
34. Takemura, Y. *et al.* Cholinergic regulation of epithelial sodium channels in rat alveolar type 2 epithelial cells. *Am J Physiol Lung Cell Mol Physiol* **304**, L428–437 (2013).
35. Bingle, L. *et al.* SPLUNC1 (PLUNC) is expressed in glandular tissues of the respiratory tract and in lung tumours with a glandular phenotype. *The Journal of pathology* **205**, 491–497 (2005).
36. Di, Y. P. *et al.* Molecular cloning and characterization of spurt, a human novel gene that is retinoic acid-inducible and encodes a secretory protein specific in upper respiratory tracts. *J Biol Chem* **278**, 1165–1173 (2003).
37. Adriaensens, D., Brouns, I., Van Genechten, J. & Timmermans, J. P. Functional morphology of pulmonary neuroepithelial bodies: extremely complex airway receptors. *Anat Rec A Discov Mol Cell Evol Biol* **270**, 25–40 (2003).
38. Jakupaj, M. *et al.* Role of endogenous NO in modulating airway contraction mediated by muscarinic receptors during development. *The American journal of physiology* **273**, L531–536 (1997).
39. Mak, J. C., Baraniuk, J. N. & Barnes, P. J. Localization of muscarinic receptor subtype mRNAs in human lung. *American journal of respiratory cell and molecular biology* **7**, 344–348 (1992).
40. Wine, J. J. Parasympathetic control of airway submucosal glands: central reflexes and the airway intrinsic nervous system. *Auton Neurosci* **133**, 35–54 (2007).
41. Restrepo, R. D. Inhaled adrenergics and anticholinergics in obstructive lung disease: do they enhance mucociliary clearance? *Respiratory care* **52**, 1159–1173; discussion 1173–1155 (2007).
42. Hirsh, A. J. *et al.* Evaluation of second generation amiloride analogs as therapy for cystic fibrosis lung disease. *The Journal of pharmacology and experimental therapeutics* **311**, 929–938 (2004).

Acknowledgements

We thank the Stanford lung transplant team, Jin Wu, Marlene Wine, Kim Tran, Tony Nguyen, Sara Modlin, Jessica Char, Rohan Verma, Marlene Wolfe, Lesje Atkinson, Miesha Farahmand, Caitlyn Lahousse, Jeeyeon Kim, Miriam Figueira, patients and their families for help with our studies. We are also grateful to Jennifer Lyons, Rosemary Broome, Armando Navarro, Bryan Blazek, Marcus Muro, Rich Henry, Zachary Sellers, Il-Ho Park, and Jin Hyeok Jeong for useful discussions and animal airway tissue supplies and Philip Karp, Michael Welsh, Jonathan Widdicombe and Walter Finkbeiner for supply human primary cell cultures. This study was supported by JOO06IO and JOO11GO (to NSJ) from Cystic Fibrosis Foundation (<http://www.cff.org>); WINE07XX0 (to JJW) from Cystic Fibrosis Foundation Therapeutics (<http://www.cff.org>); WINE13GO (to JJW) from Cystic Fibrosis Foundation (<http://www.cff.org>).

Author Contributions

N.S.J. conceived the research; N.S.J., M.E.K. and J.J.W. designed the research. N.S.J., M.E.K., J.Y.C. and H.-J.C. performed the research. N.S.J., M.E.K., J.Y.C., H.-J.C. and J.J.W. analyzed the data. N.S.J., M.E.K., J.Y.C., H.-J.C. and J.J.W. wrote the manuscript. All authors reviewed and edited the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Joo, N. S. *et al.* Inhibition of Airway Surface Fluid Absorption by Cholinergic Stimulation. *Sci. Rep.* **6**, 20735; doi: 10.1038/srep20735 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>